



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

410. Proposed by A. H. HOLMES, Brunswick, Me.

Given a focus and two tangents to an ellipse, prove that the locus of the foot of the normal corresponding to either tangent is a straight line.

No solution has been received.

411. Proposed by C. N. SCHMALL, New York City.

$ABCD$ is a rectangle of known sides. BC being produced indefinitely, it is required to draw a straight line from A cutting CD and BC in X and Y , respectively, so that the intercept XY may be equal to a given straight line. (Unsolved in *Educational Times*.)

REMARK BY L. S. SHIVELY, Mt. Morris, Ill.

This problem cannot be solved with straight edge and compasses only, for an arbitrary length of the given straight line. To prove this, it will be sufficient to show that for a particular length the solution, with these restrictions, is impossible.

Let the length of the given line be twice the diagonal of the rectangle. Suppose the problem to have been solved. Let o be the midpoint of xy . Draw AC and co . Then triangles coy and aco are isosceles and we have

$$\angle CAO = \angle AOC = 2 \angle OYC$$

and

$$\angle ACB = \angle CAO + \angle OYC = 3 \angle OYC.$$

Since the given rectangle is arbitrary we have here a solution for the trisection of an arbitrary acute angle, which is known to be impossible with straight edge and compasses. Hence the proposed construction is also impossible.

Solutions were received, too late for credit in the December number, of 403, from W. H. Johnson; of 404, from S. Lefschetz; of 406, 407, and 408 from M. A. Muzzy and H. E. Trefethen.

CALCULUS.

330. Proposed by C. N. SCHMALL, New York City.

X is a homogeneous function, in the n th degree, of x, y, z ; Y is any function of u, v, w . If $ux = vy = wz = 1 + X \dots (1)$, prove that (by Eurler's Theorem),

$$x \frac{\partial Y}{\partial x} + y \frac{\partial Y}{\partial y} + z \frac{\partial Y}{\partial z} + \frac{n}{x} \frac{\partial Y}{\partial u} + \frac{n}{y} \frac{\partial Y}{\partial v} + \frac{n}{z} \frac{\partial Y}{\partial w} = (n-1) \left(u \frac{\partial Y}{\partial u} + v \frac{\partial Y}{\partial v} + w \frac{\partial Y}{\partial w} \right).$$

SOLUTION BY THE PROPOSER.

From (1) we have

$$\log x + \log u = \log y + \log v = \log z + \log w = \log (1 + X)$$

Differentiating, we have

$$\frac{1}{x} + \frac{1}{u} \frac{\partial u}{\partial x} = \frac{1}{1+X} \frac{\partial X}{\partial x},$$

or

and similarly,

$$\left. \begin{aligned} \frac{x}{u} \frac{\partial u}{\partial x} &= \frac{x}{1+X} \frac{\partial X}{\partial x} - 1 \\ \frac{y}{u} \frac{\partial u}{\partial y} &= \frac{y}{1+X} \frac{\partial X}{\partial y} \\ \frac{z}{u} \frac{\partial u}{\partial z} &= \frac{z}{1+X} \frac{\partial X}{\partial z} \end{aligned} \right\} \quad (2)$$

We get similar expressions involving the partial derivatives of v and w , with respect to x , y and z .

Now

$$\begin{aligned} x \frac{\partial Y}{\partial x} + y \frac{\partial Y}{\partial y} + z \frac{\partial Y}{\partial z} + \frac{n}{x} \frac{\partial Y}{\partial u} + \frac{n}{y} \frac{\partial Y}{\partial v} + \frac{n}{z} \frac{\partial Y}{\partial w} &\equiv \frac{\partial Y}{\partial u} \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} \right) \\ &\quad + \frac{n}{x} \frac{\partial Y}{\partial u} + \frac{n}{y} \frac{\partial Y}{\partial v} + \frac{n}{z} \frac{\partial Y}{\partial w}. \end{aligned} \quad (3)$$

From (1),

$$x = \frac{1+X}{u}, \quad y = \frac{1+X}{v}, \quad z = \frac{1+X}{w}. \quad (4)$$

Hence, the right member of (3) can be written, in the form,

$$\frac{\partial Y}{\partial u} \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} \right) + \frac{nu}{1+X} \frac{\partial Y}{\partial u} + \frac{nv}{1+X} \frac{\partial Y}{\partial v} + \frac{nw}{1+X} \frac{\partial Y}{\partial w}. \quad (5)$$

But by adding the three equations in (2) we have

$$\frac{1}{u} \left(x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} \right) = \frac{x}{1+X} \frac{\partial X}{\partial x} + \frac{y}{1+X} \frac{\partial X}{\partial y} + \frac{z}{1+X} \frac{\partial X}{\partial z} - 1. \quad (6)$$

Now by the aid of (6) the expression (5) can be written,

$$u \frac{\partial Y}{\partial u} \left(\frac{x}{1+X} \frac{\partial X}{\partial x} + \frac{y}{1+X} \frac{\partial X}{\partial y} + \frac{z}{1+X} \frac{\partial X}{\partial z} - 1 \right) + \frac{n}{1+X} \left(u \frac{\partial Y}{\partial u} + v \frac{\partial Y}{\partial v} + w \frac{\partial Y}{\partial w} \right) \quad (7)$$

$$= u \frac{\partial Y}{\partial u} \left(\frac{nX}{1+X} - 1 \right) + \frac{n}{1+X} \left(u \frac{\partial Y}{\partial u} + v \frac{\partial Y}{\partial v} + w \frac{\partial Y}{\partial w} \right) \quad (8)$$

(by applying Euler's Theorem to the first parenthesis).

$$\begin{aligned} &= u \frac{\partial Y}{\partial u} \left(\frac{nX - X - 1}{1+X} \right) + \frac{n}{1+X} \left(u \frac{\partial Y}{\partial u} + v \frac{\partial Y}{\partial v} + w \frac{\partial Y}{\partial w} \right), \\ &= \frac{1}{1+X} \left\{ u \frac{\partial Y}{\partial u} (nX - X - 1) + n \left(u \frac{\partial Y}{\partial u} + v \frac{\partial Y}{\partial v} + w \frac{\partial Y}{\partial w} \right) \right\}. \end{aligned}$$

Remark.—The right member of the result I first gave is apparently incorrect.

Also solved by Elmer Schuyler.